

PerCP anti-human CD45

Catalog # / Size: 2120125 / 25 tests
2120130 / 100 tests

Clone: HI30

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with PerCP under optimal conditions. The solution is free of unconjugated PerCP and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: IV N816

Concentration: Lot-specific



Human peripheral blood lymphocytes stained with HI30 PerCP

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP has a maximum absorption of 482 nm and a maximum emission of 675 nm.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections and formalin-fixed paraffin-embedded tissue sections⁹, inhibition of CD45 functions⁴, immunofluorescence¹¹, and Western blotting³.

It was found that the HI30 clone and the 2D1 clone can cross block each other's binding.

Application References:

- Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press. New York.
- Kishihara K, *et al.* 1993. *Cell* 74:143.
- Esser M, *et al.* 2001. *J. Virol.* 75:6173. (WB)
- Yamada T, *et al.* 2002. *J. Biol. Chem.* 277:28830.
- Nagano M, *et al.* 2007. *Blood* 110:151.
- Jiang Q, *et al.* 2008. *Blood* 112:2858. [PubMed](#)
- Morozov A, *et al.* 2010. *Clin Cancer Res.* 16:5630. [PubMed](#)
- Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
- Friedman T, *et al.* 1999. *J. Immunol.* 162:5256. (IHC)
- Oeztuerk-Winder F, *et al.* 2012. *EMBO J.* 31:3431. (FC) [PubMed](#)
- Rees LE, *et al.* 2003. *Clin. Exp. Immunol.* 134:497. (IF)
- Lee J, *et al.* 2015. *J Exp Med.* 212:385. [PubMed](#)
- Breton G, *et al.* 2015. *J Exp Med.* 212:401. [PubMed](#)
- Marquardt N, *et al.* 2015. *J Immunol.* 6:2467. [PubMed](#)

Description: CD45 is a 180-240 kD single chain type I membrane glycoprotein also known as leukocyte common antigen (LCA) and T200. It is a tyrosine phosphatase expressed on the plasma membrane of all hematopoietic cells, except erythrocytes and platelets. CD45 is a signaling molecule that regulates a variety of cellular processes including cell growth, differentiation, cell cycle, and oncogenic transformation. CD45 plays a critical role in T and B cell antigen receptor-mediated activation by dephosphorylating substrates including p56Lck, p59Fyn, and other Src family kinases. CD45 non-covalently associates with lymphocyte phosphatase-associated phosphoprotein (LPAP) on T and B lymphocytes. CD45 has been reported to bind galectin-1 and to be associated with several other cell surface antigens including CD1, CD2, CD3, and CD4.

Antigen 1. Thomas M. 1989. *Annu. Rev. Immunol.* 7:339.
References: 2. Trowbridge I, *et al.* 1994. *Annu. Rev. Immunol.*12:85.